

AD-A031 963

WISCONSIN UNIV MADISON MATHEMATICS RESEARCH CENTER

F/G 12/1

A SURVEY OF M-MATRIX CHARACTERIZATIONS. I. NONSINGULAR M-MATRIX--ETC(U)

JUL 76 R J PLEMMONS

DAAG29-75-C-0024


NL

UNCLASSIFIED

MRC-TSR-1651

1 OF 1
AD
A031963





AD A 031 963

MRC Technical Summary Report #1651

A SURVEY OF M-MATRIX CHARACTERIZATIONS I: NONSINGULAR M-MATRICES

R. J. Plemmons

**Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706**


July 1976

(Received June 15, 1976)

**Approved for public release
Distribution unlimited**

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709



UNIVERSITY OF WISCONSIN - MADISON
MATHEMATICS RESEARCH CENTER

A SURVEY OF M-MATRIX CHARACTERIZATIONS
I: NONSINGULAR M-MATRICES

R. J. Plemmons¹⁾²⁾

Technical Summary Report #1651

July 1976

ABSTRACT

The purpose of this survey is to classify systematically a widely ranging list of characterizations of nonsingular M-matrices from the economics and mathematics literatures. These characterizations are grouped together in terms of their relationships to the properties of (1) positivity of principal minors (2) inverse-positivity and splittings (3) stability and (4) semi-positivity and diagonal dominance.

A list of forty equivalent conditions is given for a square matrix A with nonpositive off-diagonal entries to be a nonsingular M-matrix. These conditions are grouped into classes in order to identify those that are equivalent for arbitrary real matrices A .

In addition, other remarks relating nonsingular M-matrices to certain complex matrices are made and the recent literature on these general topics is surveyed.

AMS (MOS) Subject Classifications: 1502, 65F10

Key Words and Phrases: M-Matrix, Positivity of Principal Minors, Inverse-positivity and Splittings, Stability, Semi-positivity and Diagonal Dominance, Iterative Methods for Systems of Linear Equations

Work Unit Numbers 2 (Matrix Theory) and 7 (Numerical Analysis)

¹⁾Permanent address: Department of Computer Science, University of Tennessee, Knoxville, Tenn. 37916.

²⁾A portion of this research was sponsored by the U. S. Army Research Office under Contract No. DAHC04-74-C-0019, and the United States Army under Contract No. DAAG29-75-C-0024.

A SURVEY OF M-MATRIX CHARACTERIZATIONS

I: NONSINGULAR M-MATRICES

R. J. Plemmons¹⁾²⁾

I. INTRODUCTION. Very often problems in the biological, physical and social sciences can be reduced to problems involving matrices which, due to certain constraints, have some special structure. One of the most common situations is where the matrix A in question has nonpositive off-diagonal and nonnegative diagonal entries, that is, A is of the type

$$A = \begin{pmatrix} a_{11} & -a_{12} & -a_{13} & \cdot & \cdot & \cdot \\ -a_{21} & a_{22} & -a_{23} & \cdot & \cdot & \cdot \\ -a_{31} & -a_{32} & a_{33} & & & \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \end{pmatrix}$$

ACCESSION 14	
PTIS	WORLD SECTION <input checked="" type="checkbox"/>
D. C.	East Section <input type="checkbox"/>
UNCLASSIFIED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist. A. M. L. and/or SPECIAL	
A	

where the a_{ij} are nonnegative. Such matrices usually occur in relation to systems of linear or nonlinear equations or eigenvalue problems in a wide variety of areas including finite difference or finite element methods for partial differential equations, input-output production and growth models in economics, linear complementarity problems in operations research and Markov processes in probability and statistics.

¹⁾Permanent address: Department of Computer Science, University of Tennessee, Knoxville, Tenn. 37916.

²⁾A portion of this research was sponsored by the U. S. Army Research Office under Contract No. DAHC04-74-C-0019, and the United States Army under Contract No. DAAG29-75-C-0024.

We adopt the traditional notation here (of Fiedler and Ptak [1962]) by letting $Z^{n,n}$ denote the set of all $n \times n$ real matrices $A = (a_{ij})$ with $a_{ij} \leq 0$ for all $i \neq j$, $1 \leq i, j \leq n$.

Our purpose is to give a systematic treatment of the characterizations of a class of matrices in $Z^{n,n}$ first studied systematically by Ostrowski [1937].

Definition. An $n \times n$ matrix A that can be expressed in the form $A = sI - B$ where $B = (b_{ij})$ with $b_{ij} \geq 0$, $1 \leq i, j \leq n$, and $s \geq \rho(B)$, the maximum of the moduli of the eigenvalues of B , is called an M-matrix.

We shall be primarily concerned in this survey with nonsingular M-matrices and it is easy to see that this is the class of those A given in the Definition for which $s > \rho(B)$.

It should be mentioned that the theory of nonnegative matrices developed by Perron [1907a, b] and Frobenius [1908, 1909, 1912] provided a certain essential background for Ostrowski's work and the work of others on M-matrices. For example, it is easy to see from the Perron-Frobenius theory of nonnegative matrices that if A is a nonsingular M-matrix then the diagonal elements a_{ii} of A must be positive. Also, for any representation $A = sI - B$, $B \geq 0$ it follows that $s > \rho(B)$. In particular then, the class of nonsingular M-matrices forms a proper subclass of the class of M-matrices which forms a proper subclass of matrices in $Z^{n,n}$.

No attempt will be made here to systematically trace the history of the development of the theory of M-matrices. It appears however, that the term M-matrix was first used by Ostrowski [1937, 1956] in reference to the work of Minkowski [1900, 1907] who proved that if $A \in \mathbb{Z}^{n,n}$ has all of its row sums positive, then the determinant of A is positive. Papers following the early work of Ostrowski have primarily been produced by two groups of researchers, one in mathematics, the other in economics.

The mathematicians have mainly had in mind the applications of M-matrices to the establishment of bounds on eigenvalues and on the establishment of convergence criteria for iterative methods for the solution of large sparse systems of linear equations. Meanwhile, the economists have studied M-matrices in connection with gross substitutability, stability of a general equilibrium and Leontief's input-output analysis in economic systems (Leontief [1941]). With this in mind it should be mentioned that the terms productive matrix, Leontief matrix and Minkowski matrix have all been used in the economics literature to describe what we call a nonsingular M-matrix. Some of the contributions of individual researchers in these areas will be mentioned later in the paper.

Because the concept of an M-matrix has been applied to such diverse areas of the sciences, much of the theory has been developed at least partially in isolation. As a result much of the terminology and notation have been different and there is often formal overlap in the literature (this

is pointed out quite vividly by Varga [1976])). One of the purposes of this survey is to collect the various characterizations of nonsingular M-matrices that have been developed and applied by various researchers into a list that might make it possible for the reader to get an overall picture of the theory. The first systematic effort to characterize M-matrices was by Fiedler and Ptak [1962]. An initial survey of the theory of M-matrices was made by Poole and Boullion [1974]. Varga [1976] has surveyed the role of diagonal dominance in the theory of M-matrices. In addition, Schröder [1976] has surveyed some of the properties of nonsingular M-matrices using operator theory and partially ordered linear spaces. Meanwhile, Kaneko [1976] has compiled a list of characterizations and applications of nonsingular M-matrices in terms of linear complementarity problems in Operations Research. However, our objectives and our approach are quite different in this survey.

Section II is devoted to characterizing in a systematic way those matrices in $Z^{n,n}$ that are nonsingular M-matrices. Characterizations of certain real and complex matrices related to M-matrices are also given and an attempt is made to indicate where in the literature these characterizations may be found.


No proofs are given in this survey. The proofs of the various results can be found in the original papers.

The following notation will be used.

- ϕ - The empty set.
- 0 - A vector or matrix of all zeroes.
- I - An identity matrix.
- S - A signature matrix, that is, a diagonal matrix with diagonal entries $+1$ or -1 .
- \mathbb{C} - The set of complex numbers.
- \mathbb{R} - The set of real numbers.
- \mathbb{C}^n - The vector space of n -vectors over \mathbb{C} .
- \mathbb{R}^n - The vector space of n -vectors over \mathbb{R} .
- \mathbb{R}_+^n - The nonnegative orthant, that is, vectors in \mathbb{R}^n having all non-negative components. If $x \in \mathbb{R}_+^n$ we write $x \geq 0$ if $x \neq 0$, and $x \leq 0$ if $x \geq 0$ or $x = 0$. If x has all positive components we write $x > 0$.
- $\mathbb{C}^{n,n}$ - The set of all $n \times n$ matrices over \mathbb{C} .
- $\mathbb{R}^{n,n}$ - The set of $n \times n$ matrices over \mathbb{R} .
- $Z^{n,n}$ - The set of all $A = (a_{ij}) \in \mathbb{R}^{n,n}$ with $a_{ij} \leq 0$ if $i \neq j$, $1 \leq i, j \leq n$.
- Let $A \in \mathbb{R}^{n,n}$.
- $A \geq 0$ - Indicates that each component of A is nonnegative and that $A \neq 0$.
- $A \geq 0$ - Indicates that $A \geq 0$ or $A = 0$.
- A^t - The transpose of A .
- A^{-1} - The inverse of A .
- $\rho(A)$ - The spectral radius of A .

II. CHARACTERIZATIONS. For practical purposes in characterizing non-singular M-matrices, it is evident that we can often begin by assuming that $A \in Z^{n,n}$. However, many of the statements of these characterizations are equivalent without this assumption. We have attempted here to group together all such statements into certain categories. Moreover, certain other implications follow without assuming that $A \in Z^{n,n}$ and we point out such implications in the following inclusive theorem.

Theorem 1. Let $A \in \mathbb{R}^{n,n}$. Then for each fixed letter C representing one of the conditions below, conditions C_i are equivalent for each i . Moreover letting C then represent any of the equivalent conditions C_i , the following implications hold:



Finally, if $A \in Z^{n,n}$ then each of the following conditions is equivalent to the statement: A is a nonsingular M-matrix.

Positivity Of Principal Minors

A_1 . All the principal minors of A are positive.

A_2 . Every real eigenvalue of each principal submatrix of A is positive.

A₃. $A + D$ is nonsingular for each positive diagonal matrix D .

A₄. For each $x \neq 0$ there exists a positive diagonal matrix D such that

$$x^t A D x > 0 .$$

A₅. For each $x \neq 0$ there exists a nonnegative diagonal matrix D such that

$$x^t A D x > 0 .$$

A₆. A does not reverse the sign of any vector. That is, if $x \neq 0$ then for some subscript i ,

$$x_i (A x)_i > 0 .$$

A₇. For each signature matrix S there exists an $x > 0$ such that

$$S A S x > 0 .$$

B₈. The sum of all the $k \times k$ principal minors of A is positive for $k = 1, 2, \dots, n$.

C₉. Every real eigenvalue of A is positive.

C₁₀. $A + \alpha I$ is nonsingular for each scalar $\alpha \geq 0$.

D₁₁. All the leading principal minors of A are positive.

D₁₂. There exist lower and upper triangular matrices L and U respectively, with positive diagonals such that

$$A = LU .$$

E₁₃. There exists a strictly increasing sequence of subsets $\emptyset \neq S_1 \subset \dots \subset S_n = \{1, \dots, n\}$ such that the determinant of the principal submatrix of A formed by choosing row and column indices from S_i is positive for $i = 1, \dots, n$.

E_{14} . There exists a permutation matrix P and lower and upper triangular matrices L and U respectively, with positive diagonals, such that

$$PAP^t = LU.$$

Inverse-Positivity And Splittings

F_{15} . A is inverse-positive. That is, A^{-1} exists and

$$A^{-1} \geq 0.$$

F_{16} . A is monotone. That is

$$Ax \geq 0 \implies x \geq 0 \text{ for all } x \in \mathbb{R}^n.$$

F_{17} . A has a convergent regular splitting. That is, A has a representation

$$A = M - N, \quad M^{-1} \geq 0, \quad N \geq 0$$

with $M^{-1}N$ convergent. That is, $\rho(M^{-1}N) < 1$.

F_{18} . A has a convergent weak regular splitting. That is, A has a representation

$$A = M - N, \quad M^{-1} \geq 0, \quad M^{-1}N \geq 0$$

with $M^{-1}N$ convergent.

F_{19} . A has a weak regular splitting and there exists $x > 0$ with $Ax > 0$.

F_{20} . There exist inverse-positive matrices M_1 and M_2 with

$$M_1 \leq A \leq M_2.$$

F_{21} . There exists an inverse-positive matrix M , with $M \geq A$, and a nonsingular M -matrix B such that $A = MB$.

F_{22} . There exists an inverse-positive matrix M and a nonsingular M -matrix B such that $A = MB$.

G_{23} . Every weak regular splitting of A is convergent.

H_{24} . Every regular splitting of A is convergent.

Stability

I_{25} . There exists a positive diagonal matrix D such that

$$AD + DA^t$$

is positive definite.

I_{26} . A is diagonally similar to a matrix whose symmetric part is positive definite. That is, there exists a positive diagonal matrix E such that for $B = E^{-1}AE$, the matrix

$$(B + B^t)/2$$

is positive definite.

I_{27} . For each nonzero positive semi-definite matrix P , the matrix PA has a positive diagonal element.

I_{28} . Every principal submatrix of A satisfies condition I_{25} .

J_{29} . A is positive stable. That is, the real part of each eigenvalue of A is positive.

J₃₀. There exists a symmetric positive definite matrix W such that

$$AW + WA^t$$

is positive definite.

J₃₁. $A + I$ is nonsingular and

$$G = (A + I)^{-1}(A - I)$$

is convergent.

J₃₂. $A + I$ is nonsingular and for

$$G = (A + I)^{-1}(A - I),$$

there exists a positive definite symmetric matrix W such that

$$W - G^t W G$$

is positive definite.

Semi-Positivity And Diagonal Dominance

K₃₃. A is semi-positive. That is, there exists $x > 0$ with $Ax > 0$.

K₃₄. There exists $x \geq 0$ with $Ax > 0$.

K₃₅. There exists a positive diagonal matrix D such that AD has all positive row sums.

L₃₆. There exists $x > 0$ with $Ax \geq 0$ such that if $(Ax)_{i_0} = 0$, then there exist indices $1 \leq i_1, \dots, i_r \leq n$, with $a_{i_k i_{k+1}} \neq 0$ for $0 \leq k \leq r-1$ and $(Ax)_{i_r} > 0$.

M₃₇. There exists $x > 0$ with $Ax \geq 0$ and

$$\sum_{j=1}^i a_{ij} x_j > 0$$

for each $i = 1, 2, \dots, n$.

N_{38} . There exists $x > 0$ such that for each signature matrix S ,

$$SASx > 0.$$

N_{39} . A has all positive diagonal elements and there exists a positive diagonal matrix D such that AD is strictly diagonally dominant. That is,

$$a_{ii}d_i > \sum_{j \neq i} |a_{ij}|d_j$$

for $i = 1, 2, \dots, n$.

N_{40} . A has all positive diagonal elements and there exists a positive diagonal matrix D such that $D^{-1}AD$ is strictly diagonally dominant.

We remark again that Theorem 1 identifies those conditions characterizing nonsingular M -matrices that are equivalent for an arbitrary matrix in $\mathbb{R}^{n,n}$, as well as certain implications that hold for various classes of conditions. For example, A_1 through A_7 are equivalent and $A \implies B$ for an arbitrary matrix in $\mathbb{R}^{n,n}$. We remark also that it follows from the work of Schneider [1953, 1965] that a matrix $A \in \mathbb{Z}^{n,n}$ is a nonsingular M -matrix if and only if each irreducible principal submatrix of A is a nonsingular M -matrix and thus satisfies one of the equivalent conditions in Theorem 1. It should also be pointed out that some of the classes have left-right duals with A replaced by A^t .

The problem of giving proper credit to those originally responsible for the various characterizations listed in Theorem 1 is difficult, if not

impossible. The situation is complicated by the fact that many of the characterizations are implicit in the work of Perron [1907a, b] and Frobenius [1908, 1909, 1912] and in the work of Ostrowski [1937, 1956], but were not given there in their present form. Another complicating factor is that the diversification of the applications of M-matrices has led to certain conditions being derived independently. We attempt in this survey only to give references to the literature where the various proofs can be found.

First of all Condition A_1 , which is known as the Hawkins-Simon [1949] condition in the economics literature, was taken by Ostrowski [1937] as the definition for $A \in Z^{n,n}$ to be a nonsingular M-matrix. He then proceeded to show the equivalence of his definition with ours; namely that A has a representation $A = sI - B$, $B \geq 0$ and $s > \rho(B)$. Condition A_2 is also in Ostrowski [1937]. Condition A_3 was shown to be equivalent to A_1 in Wilson [1971], while Conditions A_4 , A_5 and A_6 were listed in Fiedler and Ptak [1962]. Condition A_6 is also in Gale and Nikaido [1965] and A_7 was shown to be equivalent to A_1 in Moylan [1976].

Next, Condition B_8 can be found in Johnson [1974] and C_9 , C_{10} , D_{11} , D_{12} , E_{13} and E_{14} are in Fiedler and Ptak [1962].

Condition F_{15} is in the original paper by Ostrowski [1937], Condition F_{16} was shown to be equivalent to F_{15} by Collatz [1952], Condition F_{17} is implicit in the work of Varga [1962] on regular splittings, Condition F_{18} is in the work of Schneider [1965] while F_{19} and F_{20} are in the work of Price [1968]. Conditions F_{21} , F_{22} and G_{23} are essentially in the work of Schneider [1965] and H_{24} is in Varga [1962].

Next, Condition I_{25} is in the work of Tartar [1971] and of Araki [1975]. The equivalence of I_{26} , I_{27} and I_{28} to I_{25} is in Barker, Berman and Plemmons [1976]. The stability Condition J_{29} is in the work of Ostrowski [1937] while its equivalence with J_{30} is the Lyapunov [1892] theorem. The equivalence of J_{30} with J_{31} is in Taussky [1961] and the equivalence of J_{32} with J_{31} is the Stein [1952] theorem.

Conditions K_{33} , K_{34} and K_{35} are in Schneider [1953] and Fan [1958]. Condition L_{36} in a slightly different form is in Bramble and Hubbard [1964] while this particular form is in Varga [1976]. Condition M_{37} is in the work of Beauwens [1976] and N_{38} is in that of Moylan [1976]. Conditions N_{39} and N_{40} are essentially in Schneider [1965].

Now let C represent any of the equivalent Conditions C_i and let $A \in \mathbb{R}^{n,n}$. That $N \Rightarrow I$ was established in Barker, Berman and Plemmons [1976]. That $I \Rightarrow J$ is in Lyapunov [1892] and that $I \Rightarrow A$ is also in Barker, Berman and Plemmons [1976]. The implications $A \Rightarrow B \Rightarrow C$, $A \Rightarrow D \Rightarrow E$ and $J \Rightarrow C$ are immediate. That $A \Rightarrow K$ can be found in Nikaido [1968] and the implication $K \Rightarrow L$ is immediate. That $F \Rightarrow K$ can be found in Schneider [1953]. Finally, the implication $F \Rightarrow G$ is in Varga [1962] and the implication $G \Rightarrow H$ is immediate.

It is not known by the author whether $H \Rightarrow G$. The implication will follow if it can be shown that if $A \in \mathbb{R}^{n,n}$ has a weak regular splitting then it must also have some regular splitting. It is also not known by the author whether $M \Rightarrow L$.

Next, we consider necessary and sufficient conditions for an arbitrary matrix $A \in \mathbb{R}^{n,n}$ to be a nonsingular M-matrix. In the following theorem, we do not assume that A has nonpositive off-diagonal entries.

Theorem 2. Let $A \in \mathbb{R}^{n,n}$, $n \geq 2$. Then each of the following conditions is equivalent to the statement: A is a nonsingular M-matrix.

1. $A + D$ is inverse-positive for each nonnegative diagonal matrix D .
2. $A + \alpha I$ is inverse-positive for each scalar $\alpha \geq 0$.
3. Each principal submatrix of A is inverse-positive.
4. Each principal submatrix of A of orders 1, 2 and n of A is inverse-positive.

In view of Conditions F_{15} in Theorem 1, the proof of Theorem 2 reduces to that of showing that any $A \in \mathbb{R}^{n,n}$ satisfying one of these conditions can have no positive off-diagonal elements. That this is true if Conditions 1 or 2 hold is given in Wilson [1971] and if Conditions 3 or 4 hold is in Cottle and Veinott [1972].

We now relate some of the characterizations of nonsingular M-matrices to matrices over the complex field.

For $A \in \mathbb{C}^{n,n}$ we define its comparison matrix $\mathcal{M}(A) = (m_{ij}) \in \mathbb{R}^{n,n}$ given by

$$m_{ii} = |a_{ii}|, \quad m_{ij} = -|a_{ij}|, \quad i \neq j, \quad 1 \leq i, j \leq n$$

and we define

$$\Omega(A) = \{B = (b_{ij}) \in \mathbb{C}^{n,n} : |b_{ij}| = |a_{ij}|, \quad 1 \leq i, j \leq n\},$$

to be the set of equimodular matrices associated with A .

Our objective is to characterize matrices A for which $\mathcal{M}(A)$ is a nonsingular M-matrix. Such matrices will be called H-matrices after Ostrowski [1937]. We first use Theorem 1 to characterize certain real H-matrices.

Theorem 3. Let $A \in \mathbb{R}^{n,n}$ have all positive diagonal elements. Then $\mathcal{M}(A)$ is an M-matrix if and only if A satisfies one of the equivalent Conditions N_{38} , N_{39} or N_{40} of Theorem 1.

That N_{38} is equivalent to $\mathcal{M}(A)$ being a nonsingular M-matrix is in Moylan [1976], while N_{39} and N_{40} are in Schneider [1953].

In order to characterize complex H-matrices, we introduce the following splittings of $A \in \mathbb{C}^{n,n}$. Suppose the diagonals of A are all nonzero and let

$$A = M - N = D - L - U$$

where $D = \text{diag}(a_{11}, \dots, a_{nn})$ and where $-L$ and $-U$ are respectively the lower and upper parts of A . Different choices of M in this splitting lead to certain well known iteration matrices $T = M^{-1}N$. For $\omega > 0$ we define

$$J_{\omega}(A) = \omega D^{-1}(L + U) + (1 - \omega)I$$

$$J_{\omega}^{\prime}(A) = (D - \omega L)^{-1}[(1 - \omega)D + \omega U]$$

$$J_{\omega}^{\prime\prime}(A) = (D - \omega U)^{-1}[(1 - \omega)D + \omega L](D - \omega L)^{-1}[(1 - \omega)D + \omega U]$$

respectively the (point) Jacobi overrelaxation iteration matrix, the successive overrelaxation iteration matrix and the symmetric successive overrelaxation

iteration matrix associated with A . These matrices from the iteration matrices for the widely used JOR, SOR and SSOR iteration procedures for solving systems of linear equations. We are interested in conditions under which these iteration matrices are convergent. It turns out that the theory of M-matrices plays a fundamental role in such investigations.

Our final characterization theorem is for nonsingular H-matrices in $C^{n,n}$.

Theorem 4. Let $A \in C^{n,n}$ have all nonzero diagonals. Then each of the following conditions is equivalent to the statement: A is a nonsingular H-matrix.

1. For each $B \in C^{n,n}$, $\mathcal{M}(B) \geq \mathcal{M}(A)$ implies B is nonsingular.
2. For each $B \in \Omega(A)$,

$$0 < \omega < \frac{2}{1 + \rho(J_1(B))}$$

implies that

$$\rho(J_\omega(B)) < 1.$$

3. For each $B \in \Omega(A)$,

$$0 < \omega < \frac{2}{1 + \rho(J_1(B))}$$

implies that

$$\rho(\mathcal{L}_\omega(B)) < 1.$$

4. For each $B \in \Omega(A)$,

$$0 < \omega < \frac{2}{1 + \rho(J_1(B))}$$

implies that

$$\rho(\mathfrak{S}_\omega(B)) < 1.$$

Conditions 2 and 3 in Theorem 4 are in the work of Varga [1976] in relation to characterizations of diagonal dominance and Condition 4 is in Alefeld and Varga [1976]. Finally, Condition 1 is given in Ostrowski [1956]. A related result is in Camion and Hoffman [1966], who used the theory of the alternative to show that for each $B \in \mathbb{C}^{n,n}$, $\mathfrak{M}(B) = \mathfrak{M}(A)$ implies B is nonsingular if and only if there exists a permutation matrix P such that $\mathfrak{M}(PA)$ is an M-matrix.

Acknowledgements. The author wishes to acknowledge very helpful correspondence and comments from Professors Maroslav Fiedler, Charles Johnson, Ken Kaneko, Hans Schneider and Richard Varga during the preparation of this survey.

REFERENCES

- G. Alefeld and R. S. Varga [1976], "Zur Konvergenz des Symmetrischen Relaxationsverfahrens", Numer. Math. 25, pp. 291-296.
- M. Araki [1975], "Application of M-matrices to the Stability Problems of Composite Dynamical Systems", J. Math. Anal. and Appl. 52, pp. 309-321.
- G. P. Barker, A. Berman and R. J. Plemmons [1976], "Positive Diagonal Solutions to the Lyapunov Equations", preprint.
- R. Beauwens [1976], "Semi-strict Diagonal Dominance", SIAM J. Numer. Anal. 13, pp. 104-112.
- J. H. Bramble and B. E. Hubbard [1964], "On a Finite Difference Analogue of an Elliptic Boundary Value Problem Which is Neither Diagonally Dominant Nor of Nonnegative Type", J. Math. and Phys. 43, pp. 117-132.
- P. Camion and A. J. Hoffman [1966], "On the Nonsingularity of Complex Matrices", Pacific J. Math. 17, pp. 211-214.
- L. Collatz [1952], "Aufgaben Monotoner Art", Arch. Math. 3, pp. 366-376.
- R. W. Cottle and A. F. Veinott, Jr. [1972], "Polyhedral Sets Having a Least Element", Math. Programming 3, pp. 238-249.
- Ky Fan [1958], "Topological Proofs for Certain Theorems on Matrices with Nonnegative Elements", Monatsh Math. 62, pp. 219-237.
- M. Fiedler and V. Ptak [1962], "On Matrices with Non-positive Off-diagonal Elements and Positive Principal Minors", Czechoslovak Math. J. 12, pp. 382-400.

- G. Frobenius [1908], "Über Matrizen aus Positiven Elementen",
Sitzungsberichte der Königlich Preussischen Akademie der
Wissenschaften zu Berlin, pp. 471-476.
- G. Frobenius [1909], "Über Matrizen aus Positiven Elementen, II", *ibid*,
pp. 514-518.
- G. Frobenius [1912], "Über Matrizen aus Nicht Negativen Elementen", *ibid*,
pp. 456-477.
- D. Gale and H. Nikado [1965], "The Jacobian Matrix and Global
Univalence of Mappings", Math. Annalen 19, pp. 81-93.
- D. Hawkins and H. A. Simon [1949], "Note: Some Conditions of Macro-
economic Stability", Econometrica 17, pp. 245-248.
- C. R. Johnson [1974], "A Sufficient Condition for Matrix Stability", J. Res.
National Bureau of Standards 78B, pp. 103-104.
- K. Kaneko [1976], "Linear Complementarity Problems and Characterizations
of Minkowski Matrices", preprint.
- W. W. Leontief [1941], The Structure of the American Economy, Harvard
Press, Cambridge, Mass.
- A. Lyapunov [1892], "Problèmes Général de la Stabilité du Mouvement",
Communications of the Math. Soc. of Kharkow, republished in Annals
of Math. Studies, No. 17, Princeton Press, Princeton, 1947.
- H. Minkowski [1900], "Zur Theorie der Einkerten in der Algebraischen
Zahlkörpern", Nachrichten von der Königlischen Gesellschaft der
Wissenschaften zu Gottigen, Mathematische-Physikalische Klasse,
pp. 90-93.

- H. Minkowski [1907], Diophantische Approximationen, B. G. Teubner, Leipzig, Germany.
- P. J. Moyla [1976], "Matrices with Positive Principal Minors", Linear Algebra and its Applications (to appear).
- H. Nikaido [1968], Convex Structures and Economic Theory, Academic Press, New York.
- A. Ostrowski [1937], "Über die Determinanten mit Überwiegender Hauptdiagonal", Commentarii Mathematicae Helvetici 10, pp. 69-96.
- A. Ostrowski [1956], "Determinanten mit Überwiegender Hauptdiagonal und die Absolute Konvergenz von Linearen Iterationsprozessen", ibid, 30, pp. 175-210.
- O. Perron [1907a], "Grundlagen für eine Theorie des Jacobischen Kettenbruchalgorithmus", Math. Ann. 64, pp. 1-76.
- O. Perron [1907b], "Zur Theorie der Matrizen", Math. Ann. 64, pp. 248-263.
- G. Poole and T. Boullion [1974], "A Survey on M-matrices", SIAM Review 16, pp. 419-427.
- H. S. Price [1968], "Monotone and Oscillation Matrices Applied to Finite Difference Approximation", Math. Computation 22, pp. 484-516.
- H. Schneider [1953], "An Inequality for Latent Roots Applied to Determinants with Dominant Principal Diagonal, J. London Math. Soc. 28, pp. 8-20.
- H. Schneider [1956], "The Elementary Divisors Associated with 0, of a Singular M-matrix", Proc. Edinburgh Math. Soc. 10, pp. 108-122.

- H. Schneider [1965], "Positive Operators and an Inertia Theorem", Numer. Math. 7, pp. 11-17.
- J. Schröder [1976], "M-Matrices and Generalizations", preprint.
- P. Stein [1952], "Some General Theorems on Iterants", J. Res. Natl. Bur. Standards 48, pp. 82-83.
- L. Tartar [1971], "Une Nouvelle Characterization des M-Matrices", Revue Francaise d'Informatique Rech. Oper. 5, pp. 127-128.
- O. Taussky [1961], "A Generalization of a Theorem of Lyapunov", J. Soc. Indust. Appl. Math. 9, pp. 640-643.
- O. Taussky [1964], "Matrices C with $C^n \rightarrow 0$ ", J. of Algebra 1, pp. 5-10.
- R. S. Varga [1962], "Matrix Iterative Analysis", Prentice-Hall, Englewood Cliffs, N. J.
- R. S. Varga [1976], "On Recurring Theorems on Diagonal Dominance", Linear Algebra and Its Appl. 13, pp. 1-9.
- A. N. Wilson [1971], "A Useful Generalization of the P_0 Matrix Concept", Numer. Math. 7, pp. 62-70.
- A. N. Wilson [1973], "A Note on Pairs of Matrices and Matrices of Monotone Kind", SIAM J. Numer. Anal. 10, pp. 618-622.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 1651	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER (9) Technical
4. TITLE (and Subtitle) A SURVEY OF M-MATRIX CHARACTERIZATIONS. I. NONSINGULAR M-MATRICES.		5. TYPE OF REPORT & PERIOD COVERED Summary Report, no specific reporting period
7. AUTHOR(s) R. J. Plemmons		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706		8. CONTRACT OR GRANT NUMBER(s) DAAG29-75-C-0024 DAH04-74-C-0019
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P.O. Box 12211 Research Triangle Park, North Carolina 27709		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS (12)
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) (14) MRC-TSR-1651		13. REPORT DATE (11) July 1976 12. NUMBER OF PAGES 21
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		15. SECURITY CLASS. (of this report) UNCLASSIFIED
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) M-Matrix, Stability, Positivity of Principal Minors, Inverse-positivity and Splittings, Semipositivity and Diagonal Dominance, Iterative Methods for Systems of Linear Equations		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The purpose of this survey is to classify systematically a widely ranging list of characterizations of nonsingular M-matrices from the economics and mathematics literatures. These characterizations are grouped together in terms of their relationships to the properties of (1) positivity of principal minors (2) inverse-positivity and splittings (3) stability and (4) semi-positivity and		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

221200

Next
JP

20. ABSTRACT - Cont'd.

diagonal dominance.

A list of forty equivalent conditions is given for a square matrix A with nonpositive off-diagonal entries to be a nonsingular M-matrix. These conditions are grouped into classes in order to identify those that are equivalent for arbitrary real matrices A .

In addition, other remarks relating nonsingular M-matrices to certain complex matrices are made and the recent literature on these general topics is surveyed.